

--	--	--	--	--	--	--	--	--	--

**First Semester B.E. Degree Examination, Dec.2013/Jan.2014**  
**Engineering Mathematics – I**

Time: 3 hrs.

Max. Marks:100

- Note: 1. Answer any FIVE full questions, choosing at least two from each part.**  
**2. Answer all objective type questions only in OMR sheet page 5 of the answer booklet.**  
**3. Answer to objective type questions on sheets other than OMR will not be valued.**

**PART – A**

1 a. Choose the correct answers for the following :

i) If  $y = \frac{x+2}{x+1}$ , then  $y_n$  is

A)  $\frac{(-1)^n(n+1)!}{(x+1)^{n+1}}$       B)  $\frac{(-1)^n n!}{(x+1)^{n+1}}$       C)  $\frac{(-1)^n n!}{(x+1)^n}$       D)  $\frac{(-1)^{n-1} n!}{(x+1)^{n+1}}$

ii) If  $y = (ax+b)^m$  with  $m = n$ , then  $y_n$  is

A)  $n! a^n$       B) 0      C)  $n! b^n$       D)  $n!$

iii) The geometrical interpretation of Lagrange's mean value theorem is

A)  $f'(c) = \frac{f(b)-f(a)}{b-a}$       B)  $f'(c) = \frac{f(b)+f(a)}{b-a}$       C)  $\frac{f'(c)}{g'(c)} = \frac{f(b)-f(a)}{g(b)-g(a)}$       D) none of these

iv) The Maclaurin's series expansion of  $e^{-x}$  is

A)  $1+x+\frac{x^2}{2!}+\frac{x^3}{3!}+\dots$       B)  $1-x+\frac{x^2}{2!}-\frac{x^3}{3!}+\dots$   
 C)  $x-\frac{x^2}{2!}+\frac{x^3}{3!}-\dots$       D)  $x+\frac{x^2}{2!}+\frac{x^3}{3!}+\dots$       (04 Marks)

b. If  $y = \sin \log (x^2 + 2x + 1)$ , prove that  $(x+1)^2 y_{n+2} + (2n+1)(x+1)y_n + (n^2+4)y_n = 0$ .

(04 Marks)

c. If  $x$  is positive, show that  $x > \log(1+x) > x - \frac{1}{2}x^2$ .

(06 Marks)

d. Using Maclourin's series, expand  $\log(1+e^x)$  upto the terms containing  $x^4$ .

(06 Marks)

2 a. Choose the correct answers for the following :

i)  $\lim_{x \rightarrow \frac{\pi}{4}} \left( \frac{1 - \tan x}{\frac{\pi}{4} - x} \right)$  is equal to

A) 2      B) -2      C) 1      D) -1

ii) If  $\phi$  be the angle between the tangent and radius vector at any point on the curve  $r = f(\theta)$ , then  $\sin \phi$  is equal to

A)  $dr/ds$       B)  $r \frac{d\theta}{ds}$       C)  $r \frac{dr}{d\theta}$       D)  $ds/dr$

iii) The rate at which the curve is bending called

A) radius of curvature      B) curvature      C) circle of curvature      D) evolute

iv) The radius of curvature for polar curve  $r = f(\theta)$  is given by

A)  $\frac{(r^2 + r_1^2)^{3/2}}{r^2 + r_1^2 - rr_2}$       B)  $\frac{(r^2 + r_1^2)^{3/2}}{r_1^2 + 2r^2 - rr_2}$       C)  $\frac{(r^2 + r_1^2)^{3/2}}{r^2 + 2r_1^2 - rr_2}$       D)  $\frac{(r^2 - r_1^2)^{3/2}}{r^2 + 2r_1^2 - rr_2}$       (04 Marks)

b. Find the Pedal equation of the curve  $r^m = a^m \cos m\theta$ .

(04 Marks)

c. Find the radius of curvature for the curve  $y^2 = \frac{a^2(a-x)}{x}$ , where the curve meets the  $x$ -axis.

(06 Marks)

d. Evaluate  $\lim_{x \rightarrow \infty} \left( \frac{ax+1}{ax-1} \right)^x$ .

(06 Marks)

3 a. Choose the correct answers for the following :

i) If  $u = \log(x^2 + y^2 + z^2)$ , then  $\frac{\partial u}{\partial z}$  is

A)  $\frac{2x}{x^2 + y^2 + z^2}$       B)  $\frac{2y}{x^2 + y^2 + z^2}$       C)  $\frac{2z}{x^2 + y^2 + z^2}$       D)  $\frac{2z}{x^2 + y^2 - z^2}$

ii) If  $u = f(x, y)$  and  $y$  is a function  $x$ , then

A)  $\frac{du}{dx} = \frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} \frac{dy}{dx}$       B)  $\frac{\partial u}{\partial x} = \frac{du}{dx} + \frac{\partial u}{\partial y} \frac{dy}{dx}$   
 C)  $\frac{du}{dx} = \frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial x}$       D)  $\frac{\partial u}{\partial x} = \frac{du}{dx} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial x}$

iii) If  $r = \frac{\partial^2 f}{\partial x^2}$ ,  $s = \frac{\partial^2 f}{\partial x \partial y}$  &  $t = \frac{\partial^2 f}{\partial y^2}$ , then the condition for the saddle point is

A)  $rt - s^2 < 0$       B)  $rt - s^2 = 0$       C)  $rt - s^2 > 0$       D)  $rt - s^2 \neq 0$

iv) If  $u = x + y + z$ ,  $v = y + z$ ,  $z = z$ , then  $J\left(\frac{u, v, z}{x, y, z}\right)$  is equal to

A) 1      B) -1      C) 0      D) none of these  
 (04 Marks)

b. The focal length of a mirror is given by the formula  $\frac{1}{v} - \frac{1}{u} = \frac{2}{f}$ . If equal errors, 'e' are made

in the determination of  $u$  and  $v$ . show that the resulting error in  $f$  is  $e\left(\frac{1}{u} + \frac{1}{v}\right)$ . (04 Marks)

c. If  $u = f(2x - 3y, 3y - 4z, 4z - 2x)$ , prove that  $\frac{1}{2} \frac{\partial u}{\partial x} + \frac{1}{3} \frac{\partial u}{\partial y} + \frac{1}{4} \frac{\partial u}{\partial z} = 0$ . (06 Marks)

d. If  $x = u(1 - v)$ ,  $y = uv$ . Prove that  $JJ' = 1$ . (06 Marks)

4 a. Choose the correct answers for the following :

i) Directional derivative is maximum along

A) tangent to the surface      B) normal to the surface  
 C) any unit vector      D) coordinate axes

ii) If  $r = |x_i + y_j + 2k|$ , then  $\nabla r^n$  is

A)  $nr^{n-1}$       B)  $r^{n-1}$       C)  $\nabla \cdot \nabla r^n$       D) none of these

iii) If  $f = 3x^2 - 3y^2 + 4z^2$ , then curl (grad  $f$ ) is

A)  $4x - 6y + 8z$       B)  $4x_i - 6y_j + 8z k$       C)  $\vec{0}$       D) 3

iv) If the base vectors  $e_1$  and  $e_2$  are orthogonal then  $|e_1 \times e_2|$  is

A) 0      B) -1      C) +1      D) none of these  
 (04 Marks)

b. If  $\vec{F} = (x + y + 1)i + j - (x + y)k$ , show that  $\vec{F} \cdot \text{curl } \vec{F} = 0$ . (04 Marks)

c. Find constants 'a' and 'b' such that  $\vec{F} = (axy + z^3)i + (3x^2 - z)j + (bxz^2 - y)k$  is irrotational.

Also find a scalar function  $\phi$  such that  $\vec{F} = \nabla \phi$ . (06 Marks)

d. Prove that a spherical coordinate system is orthogonal. (06 Marks)

## PART – B

5 a. Choose the correct answers for the following :

i)  $\int_0^{\pi} \sin^7 x \, dx$  is equal to

- A) zero                      B)  $\frac{32\pi}{35}$                       C)  $\frac{32}{35}$                       D)  $= \frac{35\pi}{32}$

ii) The asymptote of  $(2 - x)y^2 = x^3$  is

- A)  $x = 2$                       B)  $y$  - axis                      C)  $x$  - axis                      D) none of these

iii) The area of the cardioid  $r = a(1 - \cos\theta)$  is

- A)  $\frac{3\pi a^2}{2}$                       B)  $\frac{3\pi}{2}$                       C)  $\frac{a^2}{2}$                       D)  $\frac{3a^2}{2}$

iv) The entire length of the asteroid  $x^{2/3} + y^{2/3} = a^{2/3}$  is

- A)  $6a$                       B)  $3a$                       C)  $2a$                       D)  $a$ .                      (04 Marks)

b. Evaluate  $\int_0^{\pi} \log(1 + a \cos x) dx$  by differentiating under the integral sign.                      (04 Marks)

c. Evaluate  $\int_0^{2a} x^2 \sqrt{2ax - x^2} dx$ , using reduction formula.                      (06 Marks)

d. Find the volume of generated by the revolution of the curve  $r = a(1 + \cos \theta)$  about the initial line.                      (06 Marks)

6 a. Choose the correct answers for the following :

i) The general solution of the differential equation  $dy/dx = (y/x) + \tan(y/x)$  is  
 A)  $\sin(y/x) = c$                       B)  $\sin(y/x) = cx$                       C)  $\cos(y/x) = cx$                       D)  $\cos(y/x) = c$

ii) The family of straight lines passing through the origin is represented by the differential equation :

- A)  $ydx + xdy = 0$                       B)  $xdy - ydx = 0$                       C)  $xdx + ydy = 0$                       D)  $ydy - xdx = 0$

iii) The homogeneous differential equation  $Mdx + Ndy = 0$  can be reduced to a differential equation, in which the variables are separated by the substitution

- A)  $y = vx$                       B)  $x + y = v$                       C)  $xy = v$                       D)  $x - y = v$

iv) The equation  $y - 2x = c$  represents the orthogonal trajectories of the family

- A)  $y = ae^{-2x}$                       B)  $x^2 + 2y^2 = a$                       C)  $xy = a$                       D)  $x + 2y = a$                       (04 Marks)

b. Solve  $(x + 1) \frac{dy}{dx} - y = e^{3x}(x + 1)^2$ .                      (04 Marks)

c. Solve  $(1 + xy) ydx + (1 - xy) xdy = 0$ .                      (06 Marks)

d. Find the orthogonal trajectory of the cardioids  $r = a(1 - \cos \theta)$ .                      (06 Marks)

- 7 a. Choose the correct answers for the following :
- If every minor of order 'r' of a matrix A is zero, then rank of A is  
A) greater than r      B) equal r      C) less than or equal to r      D) less than r.
  - The trivial solution for the given system of equations  $x + 2y + 3z = 0$ ,  $3x + 4y + 4z = 0$ ,  $7x + 10y + 12z = 0$  is  
A) (1, 1, 1)      B) (1, 0, 0)      C) (0, 1, 0)      D) (0, 0, 0)
  - Matrix has a value. This statement  
A) is always true      B) depends upon the matrices C) is false      D) none of these
  - If A is singular and  $\rho(A) = \rho(A : B)$  then the system has  
A) unique solution      B) infinitely many solution C) trivial solution D) no solution.  
(04 Marks)
- b. Using elementary transformations, find the rank of the matrix
- $$\begin{bmatrix} 1 & 2 & 0 & -1 \\ 3 & 4 & 1 & 2 \\ -2 & 3 & 2 & 5 \end{bmatrix}.$$
- (04 Marks)
- c. Show that the system  $x + y + z = 4$ ;  $2x + y - z = 1$ ;  $x - y + 2z = 2$  is consistent, solve the system.  
(06 Marks)
- d. Apply Gauss – Jordan method to solve the system of equation :  
 $2x + 5y + 7z = 52$ ;  $2x + y - z = 0$ ;  $x + y + z = 9$ .  
(06 Marks)
- 8 a. Choose the correct answers for the following :
- A square matrix A is called orthogonal, if  
A)  $A = A^L$       B)  $A^T = A^{-1}$       C)  $AA^{-1} = I$       D) none of these
  - The eigen values of the matrix  $\begin{bmatrix} 2 & \sqrt{2} \\ \sqrt{2} & 2 \end{bmatrix}$  are  
A)  $1 \pm \sqrt{6}$       B)  $1 \pm \sqrt{5}$       C)  $\sqrt{5}$       D) 1
  - The index and signature of the quadratic form  $x_1^2 + 2x_2^2 - 3x_3^2$  are respectively  
A) 2, 1      B) 1, 2      C) 1, 1      D) none of these
  - Two square matrices A and B are similar, if  
A)  $A = B$       B)  $B = P^{-1}AP$       C)  $A^T = B^T$       D)  $A^{-1} = B^{-1}$ .  
(04 Marks)
- b. Reduce the quadratic form  $8x^2 + 7y^2 + 3z^2 - 12yz + 4zx - 8xy$  to the canonical form  
(04 Marks)
- c. Determine the characteristic roots and eigen vectors of
- $$A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}.$$
- (06 Marks)
- d. Reduce the quadratic form  $x_1^2 + 2x_2^2 - 7x_3^2 - 4x_1x_2 + 8x_2x_3$  into sum of squares. (06 Marks)

\*\*\*\*\*