

## Applied maths - I

D: PH (April Exam) 181

Con. 6865-13.

(REVISED COURSE)

GS-5103

(3 Hours)

[ Total Marks : 80

N.B. (1) Question No. 1 is compulsory.

(2) Attempt any **three** questions from Question Nos. 2 to Questions No. 6(3) **Figures** to the **right** indicate **full** marks.

1. (a) If  $\cos hx = \sec \theta$  prove that  $x = \log (\sec \theta + \tan \theta)$ . 3
- (b) If  $u = \log (x^2 + y^2)$ , prove that  $\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x}$  3
- (c) If  $x = r \cos \theta$ ,  $y = r \sin \theta$ . Find  $\frac{\partial(x, y)}{\partial(r, \theta)}$ . 3
- (d) Expand  $\log (1 + x + x^2 + x^3)$  in powers of  $x$  upto  $x^8$ . 3
- (e) Show that every square matrix can be uniquely expressed as sum of a symmetric and a Skew-symmetric matrix. 4
- (f) Find  $n^{\text{th}}$  order derivative of 4  
 $y = \cos x. \cos 2x. \cos 3x.$
2. (a) Solve the equation  $x^6 - i = 0$ . 6
- (b) Reduce matrix A to normal form and find its rank where :- 6
- $$A = \begin{bmatrix} 1 & 2 & 3 & 2 \\ 2 & 3 & 5 & 1 \\ 1 & 3 & 4 & 5 \end{bmatrix}$$
- (c) State and prove Euler's theorem for a homogeneous function in two variables and 8  
 hence find  $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$  where  $u = \frac{\sqrt{x} + \sqrt{y}}{x + y}$
3. (a) Determine the values of  $\lambda$  so that the equations  $x + y + z = 1$ ;  $x + 2y + 4z = \lambda$ ;  $x + 4y + 10z = \lambda^2$  have a solution and solve them completely in each case. 6
- (b) Find the stationary values of 6  
 $x^3 + y^3 - 3axy$ ,  $a > 0$ .
- (c) Separate into real and imaginary parts  $\tan^{-1} (e^{i\theta})$ . 8

[ TURN OVER

4. (a) If  $x = u \cos v$ ,  $y = u \sin v$  6

Prove that  $\frac{\partial(x, y)}{\partial(u, v)} \cdot \frac{\partial(u, v)}{\partial(x, y)} = 1$ .

- (b) If  $\tan [\log (x + iy)] = a + ib$ , prove that  $\tan [\log (x^2 + y^2)] = \frac{2a}{(1 - a^2 - b^2)}$  where 6

$a^2 + b^2 \neq 1$ .

- (c) Using Gauss-Siedel iteration method, solve 8

$$10x_1 + x_2 + x_3 = 12$$

$$2x_1 + 10x_2 + x_3 = 13$$

$$2x_1 + 2x_2 + 10x_3 = 14$$

upto three iterations.

5. (a) Expand  $\sin^7 \theta$  in a series of sines of multiples of  $\theta$ . 6

- (b) Evaluate  $\lim_{x \rightarrow 0} \frac{(x^x - x)}{(x - 1 - \log x)}$  6

- (c) If  $y^{1/m} + y^{-1/m} = 2x$ , prove that 8

$$(x^2 - 1) y_{n+2} + (2n + 1) xy_{n+1} + (n^2 - m^2) y_n = 0.$$

6. (a) Examine the following vectors for linear dependence/Independence. 6

$$X_1 = (a, b, c), X_2 = (b, c, a), X_3 = (c, a, b) \text{ where } a + b + c \neq 0.$$

- (b) If  $z = f(x, y)$ ,  $x = e^u + e^{-v}$ ,  $y = e^{-u} - e^v$ , prove that 6

$$\frac{\partial z}{\partial u} - \frac{\partial z}{\partial v} = x \frac{\partial z}{\partial x} - y \frac{\partial z}{\partial y}$$

- (c) Fit a straight line to the following data and estimate the production in the year 1957. 8

<b>Year :</b>	1951	1961	1971	1981	1991
<b>Production in the Thousand tons :</b>	10	12	08	10	13